

AD-A072 762 BROWN UNIV PROVIDENCE R I LEFSCHETZ CENTER FOR DYNAM--ETC F/G 12/1  
ON THE EXISTENCE OF SIMPLE LIAPUNOV FUNCTIONS FOR LINEAR RETARD--ETC(U)  
JUL 79 L A CARVALHO, E F INFANTE, J A WALKER N00014-76-C-0278

UNCLASSIFIED

AFOSR-TR-79-0920

NL

| OF |  
AD  
A072762





REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR-79-0920</b>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>ON THE EXISTENCE OF SIMPLE LIAPUNOV FUNCTIONS FOR LINEAR RETARDED DIFFERENCE DIFFERENTIAL EQUATIONS.</b>		5. TYPE OF REPORT & PERIOD COVERED <b>Interim / repty</b>
7. AUTHOR(s) <b>L.A.V. Carvalho, E.F. Infante and J.A. Walker</b>		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>Brown University Division of Applied Mathematics Providence, Rhode Island 02912</b>		8. CONTRACT OR GRANT NUMBER(s) <b>AFOSR-76-3092</b>
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 20332</b>		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>61102F 2304/A1</b>
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <b>1234 P.</b>		12. REPORT DATE <b>July 1979</b>
		13. NUMBER OF PAGES <b>28</b>
		15. SECURITY CLASS. (of this report) <b>UNCLASSIFIED</b>
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  <b>Approved for public release; distribution unlimited.</b>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20; if different from Report) <b>15 N00014-76-C-0278 AFOSR-76-3092</b>		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  <b>Liapunov functions of simple form have been used for the study of stability properties of difference-differential equations. In this paper, we provide necessary and sufficient conditions for the existence of such functions.</b>		

**LEVEL**

**DDC  
RECEIVED  
AUG 14 1979  
C**

**AD A072762**

**DDC FILE COPY**

ON THE EXISTENCE OF SIMPLE LIAPUNOV FUNCTIONS FOR  
LINEAR RETARDED DIFFERENCE DIFFERENTIAL EQUATIONS

by

L.A.V. Carvalho<sup>\*</sup>  
Lefschetz Center for Dynamical Systems  
Division of Applied Mathematics  
Brown University  
Providence, R.I. 02912  
and  
Universidade Federal de Goiás  
Goiânia, Brazil, 74000

E. F. Infante<sup>\*\*</sup>  
Lefschetz Center for Dynamical Systems  
Division of Applied Mathematics  
Brown University  
Providence, R.I. 02912

J. A. Walker<sup>\*\*\*</sup>  
Lefschetz Center for Dynamical Systems  
Division of Applied Mathematics  
Brown University  
Providence, R.I. 02912  
and  
Department of Mechanical Engineering  
Northwestern University  
Evanston, Illinois 60201

July, 1979

---

<sup>\*</sup> This research was partially supported by CAPES (Coordenação do Aperfeiçoamento de Pessoal de Nível Superior - Brasil).

<sup>\*\*</sup> This research was supported in part by the Air Force Office of Scientific Research under AF-AFOSR 76-3092C, in part by the National Science Foundation under NSF-MCS 78-18858, in part by the Office of Naval Research under N00014-76-C-0278-P0002, and in part by the United States Army under ARO-D-31-124-73-G130.

<sup>\*\*\*</sup> This research was supported in part by the National Science Foundation under NSF-ENG.76-81570.



ABSTRACT

ON THE EXISTENCE OF SIMPLE LIAPUNOV  
FUNCTIONS FOR LINEAR RETARDED DIFFERENCE  
DIFFERENTIAL EQUATIONS

by

L.A.V. Carvalho, E.F. Infante, J. A. Walker

Liapunov functions of simple form have been used for the study of stability properties of difference-differential equations. In this paper we provide necessary and sufficient conditions for the existence of such functions.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF SUBMITTAL TO DDC  
This document report has been reviewed and is  
approved for public release under E.O. 13526 (7b).  
Distribution is unlimited.  
A. D. FLORE  
Technical Information Officer

Accompanying Statement

ON THE EXISTENCE OF SIMPLE LIAPUNOV  
FUNCTIONS FOR LINEAR RETARDED DIFFERENCE  
DIFFERENTIAL EQUATIONS

by

L.A.V. Carvalho, E.F. Infante, J.A. Walker

Accession For	
NTIS Serial	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Availand/or special

In this paper we consider, for the retarded linear autonomous difference-differential equation

$$\dot{y}(t) = Ay(t) + \sum_{k=1}^m B_k y(t-\tau_k), \quad t \geq 0,$$

the question of existence of Liapunov functions of the two forms

$$V(x) = y^* R y + \sum_{k=1}^m \int_{-\tau_k}^0 V^*(\theta) Q_k V(\theta) d\theta,$$

and

$$W(x) = \sup_{\max_{k=1, \dots, m} \tau_k \leq \theta \leq 0} V^*(\theta) R V(\theta).$$

It is noted that the existence of a Liapunov function of either of the foregoing forms is independent of the delays  $\{\tau_k\}_{k=1}^m$ ; hence, so will be any stability conclusions drawn from the use of such functions. This remark suggests that the class of functional equations of the above form for which such functions exist is narrow. In the paper we give a characterization of this class; moreover, we show by counterexample that there do exist equations of this form that yield asymptotic stability for all delays  $\{\tau_k\}_{k=1}^m$

and yet there exist no Liapunov function of the above forms that can be used to establish this fact.

Finally, we use Liapunov functions of the above type to provide estimates of exponential decay.

ON THE EXISTENCE OF SIMPLE LIAPUNOV  
FUNCTIONS FOR LINEAR RETARDED DIFFERENCE  
DIFFERENTIAL EQUATIONS

Dedicated to Professor Taro Yoshizawa  
on the occasion of his sixtieth birthday

by

L.A.V. Carvalho

E. F. Infante

J. A. Walker

## 1. Introduction

In this paper we consider the existence of two simple types of Liapunov functions for retarded linear autonomous difference-differential equations. The study of the stability properties of these equations was pioneered by Yoshizawa [1,2], Razumikhin [3], Krasovskii [4], Driver [5], and Hale [6].

Restriction to Liapunov functions of simple form leads to stability criteria which are overly sufficient; on the other hand, such functions are particularly desirable in specific applications where ease of computation is of primary importance. Here we provide necessary and sufficient conditions for the existence of useful Liapunov functions having standard simple forms for certain difference-differential equations.

Consider the difference-differential equation

$$\dot{y}(t) = Ay(t) + \sum_{k=1}^m B_k y(t-\tau_k), \quad t \geq 0, \quad (1.1)$$

with initial data

$$y(\theta) = v_0(\theta), \quad -\tau \leq \theta \leq 0,$$

where  $\tau_k \geq 0$ ,  $\tau \equiv \max_k \tau_k$ , and the  $n \times n$  complex matrices  $A, B_k$  are given, as is the initial function  $v_0: [-\tau, 0] \rightarrow \mathbb{C}^n$ .

Let us replace the above equation by an abstract evolution

equation on a Banach space  $\chi$ ,

$$\begin{aligned}\dot{x}(t) &= Fx(t), \quad t \geq 0, \\ x(0) = x_0 &\equiv (v_0(0), v_0) \in \mathcal{D}(F) \subset \chi,\end{aligned}\tag{1.2}$$

where  $\chi \equiv \{x = (y, v) \in \mathbb{C}^n \times \mathcal{C}([- \tau, 0]; \mathbb{C}^n) \mid y = v(0)\}$  with

$\|x\|_{\chi} \equiv \sup_{-\tau \leq \theta \leq 0} \|v(\theta)\|_{\mathbb{C}^n}$ ; the operator  $F$  has domain

$$\mathcal{D}(F) \equiv \{x = (y, v) \in \chi \mid v' \in \mathcal{C}([- \tau, 0]; \mathbb{C}^n), v'(0) = Ay + \sum_{k=1}^m B_k v(-\tau_k)\}, \text{ where}$$

$$v'(\theta) \equiv \frac{d}{d\theta} v(\theta), \text{ and is defined by } Fx \equiv (Ay + \sum_{k=1}^m B_k v(-\tau_k), v')$$

for all  $x = (y, v) \in \mathcal{D}(F)$ . It is well known [7] that  $F$  is the infinitesimal generator of a  $C_0$ -semigroup  $\{S(t)\}_{t \geq 0}$  of bounded linear operators  $S(t): \chi \rightarrow \chi$ ; in particular, for every  $x_0 \in \mathcal{D}(F)$ ,  $S(\cdot)x_0$  is the unique strong solution of (1.2).

We are interested in obtaining stability results for (1.2) (hence (1.1)) through the use of Liapunov functions.

Definition 1.1: A continuous function  $V: \chi \rightarrow \mathbb{R}$  is said to be a continuous Liapunov function for  $\{S(t)\}_{t \geq 0}$  on  $\chi$  if  $\dot{V}(x) \leq 0$  for all  $x \in \chi$ , where  $\dot{V}: \chi \rightarrow \mathbb{R}$  is given by

$$\dot{V}(x) \equiv \liminf_{t \searrow 0} \frac{1}{t} [V(S(t)x) - V(x)], \quad x \in \chi.$$



It is well known that if  $V$  is a Liapunov function on  $\chi$ , then  $V(S(t)x) \leq V(x)$  for all  $t \geq 0$ ,  $x \in \chi$ . The major difficulty encountered in using such a conclusion to derive stability results is the construction of a suitable Liapunov function.

In [8,9] very complicated Liapunov functions have been shown to yield necessary and sufficient conditions for stability and asymptotic stability for (1.2), hence for (1.1). On the other hand, most attention [4,6,7] has been centered on the use of functions having very simple structure; in particular, functions of the form

$$V(x) = y^* R y + \sum_{k=1}^m \int_{-\tau_k}^0 v^*(\theta) Q_k v(\theta) d\theta, \quad x = (y, v) \in \chi, \quad (1.3)$$

or of the form

$$W(x) = \sup_{-\tau \leq \theta \leq 0} v^*(\theta) R v(\theta), \quad x = (y, v) \in \chi, \quad (1.4)$$

where  $y^*$  denotes the conjugate transpose of the column matrix  $y$ , and  $R, Q_k$ , are  $n \times n$  Hermitian matrices. Computing  $\dot{V}: \chi \rightarrow \mathbb{R}$  and  $\dot{W}: \chi \rightarrow \mathbb{R}$  according to (1.2), we obtain

$$\begin{aligned} \dot{V}(x) = & y^* (R A + A^* R + \sum_{k=1}^m Q_k) y + 2 \operatorname{Re} \sum_{k=1}^m y^* R B_k v(-\tau_k) \\ & - \sum_{k=1}^m v^*(-\tau_k) Q_k v(-\tau_k), \end{aligned} \quad (1.5)$$

and

$$\dot{W}(x) \leq \begin{cases} 0 & \text{if } y^* R y < W(x), \\ \max\{0, 2 \operatorname{Re} y^* R (A y + \sum_{k=1}^m B_k v(-\tau_k))\} & \text{if } y^* R y = W(x), \end{cases} \quad (1.6)$$

for all  $x = (y, v) \in \chi$ ; hence,  $V$  is a Liapunov function on  $\chi$  if and only if

$$\begin{bmatrix} RA + A^* R + \sum_{k=1}^m Q_k & RB_1 & RB_2 & \dots & RB_m \\ B_1^* R & -Q_1 & 0 & \dots & 0 \\ B_2^* R & 0 & -Q_2 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ B_m^* R & 0 & 0 & \dots & -Q_m \end{bmatrix} \leq 0 \quad (1.7)$$

(i.e., this matrix is negative-semidefinite), while  $W$  is a Liapunov function on  $\chi$ , if and only if

$$\operatorname{Re} z_0^* R (A z_0 + \sum_{k=1}^m B_k z_k) \leq 0 \quad (1.8)$$

for all  $z_0, z_1, \dots, z_m \in \mathbb{C}^n$  such that  $z_0^* R z_0 \geq z_k^* R z_k$  for all  $k = 1, 2, \dots, m$ . We remark that under condition (1.8) the Liapunov

function  $W(x)$  corresponds to a "Razumikhin function" [3,7] of the form  $y^* R y$ .

It is noted that the existence of a Liapunov function of either of the foregoing forms is independent of the delays  $\{\tau_k\}_{k=1}^n$ ; hence so will be any stability conclusions drawn from the use of such functions. This remark suggests that the class of equations of the form (1.1) for which such functions exist is narrow. In the sequel we give a characterization of this class; moreover, we show by counterexample that there do exist equations of the form (1.1) that yield asymptotic stability for all delays  $\{\tau_k\}_{k=1}^m$  and yet there exists no simple Liapunov function, having either of the foregoing forms, that can be used to establish this fact.

## II. The Single Delay Case

In this section we consider the case where equation (1.2) involves only a single delay ( $m = 1$ ,  $\tau \equiv \tau_1$ ).

Essential to our arguments is the following result.

Lemma 2.1: Given  $n \times n$  complex matrices  $M, N, P$ , with  $M$  and  $N$  Hermitian, the following two statements are equivalent:

- (i)  $M + N + e^{i\phi}P + e^{-i\phi}P^* \leq 0$  for all  $\phi \in \mathbb{R}$ .
- (ii) There exists a Hermitian matrix  $Q$  such that

$$\begin{bmatrix} M + Q & P \\ P^* & N - Q \end{bmatrix} \leq 0.$$

Proof: Noting that (ii) implies

$$[I, e^{i\phi}I] \begin{bmatrix} M + Q & P \\ P^* & N - Q \end{bmatrix} \begin{bmatrix} I \\ e^{i\phi}I \end{bmatrix} = M + N + e^{i\phi}P + e^{-i\phi}P^* \leq 0$$

for all  $\phi \in \mathbb{R}$ , we see that (ii) implies (i).

To show the converse, note from (i) that if  $\hat{u}(i\phi)$  denotes the fourier transform of an arbitrary  $u(t)$ ,  $u \in \mathcal{L}_2(\mathbb{R}; \mathbb{C}^n)$ , then

$$\int_{-\infty}^{\infty} \hat{u}^*(i\phi) [M + N + e^{i\phi}P + e^{-i\phi}P^*] \hat{u}(i\phi) d\phi \leq 0; \text{ hence, by Parseval's equation,}$$

$$\int_{-\infty}^{\infty} [u^*(t)(M+N)u(t) + 2\operatorname{Re}u^*(t)Pu(t-1)]dt \leq 0.$$

Let  $u(t) = 0$  for  $t < 0$  and  $u(t) = u_k$  for  $k \leq t < k+1$ ,  $k = 0, 1, 2, \dots$ ; it then follows that

$$\sum_{k=0}^{\infty} [u_k^*(M+N)u_k + 2\operatorname{Re}u_k^*Pu_{k-1}] \leq 0$$

for all  $u_k \in \mathbb{C}^n$  such that  $\{u_k\}_{k=0}^{\infty} \in \ell_2^n$ ,  $u_{-1} \equiv 0$ .

Define the functional

$$J(u_0) \equiv \sup_{\{u_k\}_{k=1}^{\infty}} \sum_{k=1}^{\infty} [u_k^*(M+N)u_k + 2\operatorname{Re}u_k^*Pu_{k-1}], \quad u_0 \in \mathbb{C}^n.$$

We note that  $-u_0^*(M+N)u_0 \geq J(u_0) \geq 0$  and, therefore,  $J(0) = 0$  and  $J(u_0)$  is continuous at  $u_0 = 0$ ; furthermore, it is easily verified that  $J(\gamma u_0) = |\gamma|^2 J(u_0)$  for all  $\gamma \in \mathbb{C}$ .

We first wish to show that  $J^{1/2}(\cdot)$  is a seminorm on  $\mathbb{C}^n$ ; it only remains to be shown that  $J^{1/2}(\cdot)$  satisfies the triangle inequality. To this end, consider positive real numbers  $\alpha, \beta$ , such that  $\alpha^2 + \beta^2 = 1$ , and notice that, for arbitrary  $w, z \in \mathbb{C}^n$ ,

$$\begin{aligned}
J(w+z) &= \sup_{\{u_k\}_{k=1}^{\infty}} \{u_1^*(M+N)u_1 + 2\operatorname{Re} u_1^* P(w+z) + \sum_{k=2}^{\infty} [u_k^*(N+M)u_k + 2\operatorname{Re} u_k^* P u_{k-1}]\} \\
&\leq \sup_{\{u_k\}_{k=1}^{\infty}} \{(\alpha u_1)^*(M+N)(\alpha u_1) + 2\operatorname{Re}(\alpha u_1)^* P(\alpha^{-1}w) + \\
&\quad \sum_{k=2}^{\infty} [(\alpha u_k)^*(M+N)(\alpha u_k) + 2\operatorname{Re}(\alpha u_k)^* P(\alpha u_{k-1})]\} + \\
&\quad \sup_{\{u_k\}_{k=1}^{\infty}} \{(\beta u_1)^*(M+N)(\beta u_1) + 2\operatorname{Re}(\beta u_1)^* P(\beta^{-1}z) + \\
&\quad \sum_{k=2}^{\infty} [(\beta u_k)^*(M+N)(\beta u_k) + 2\operatorname{Re}(\beta u_k)^* P(\beta u_{k-1})]\} \\
&= J(\alpha^{-1}w) + J(\beta^{-1}z) = \alpha^{-2}J(w) + \beta^{-2}J(z).
\end{aligned}$$

If  $J(w) = 0 = J(z)$ , it follows that  $J(w+z) = 0$ ; if not, let  $\alpha^2 \equiv J^{1/2}(w)/[J^{1/2}(w) + J^{1/2}(z)]$ ,  $\beta^2 \equiv 1 - \alpha^2$ , and note that we obtain  $J(w+z) \leq J(w) + 2J^{1/2}(w)J^{1/2}(z) + J(z)$ . Therefore,  $J^{1/2}(\cdot)$  is a seminorm on  $\mathbb{C}^n$ .

We now wish to show that this seminorm satisfies the parallelogram law. Let  $w_0, z_0 \in \mathbb{C}^n$ ,  $\varepsilon > 0$ , and note that there exist sequences  $\{w_k\}_{k=1}^{\infty}$ ,  $\{z_k\}_{k=1}^{\infty} \in \mathbb{C}_2^n$  such that



$$J(w_0) \leq \varepsilon + \sum_{k=1}^{\infty} [w_k^* (M+N) w_k + 2\operatorname{Re} w_k^* P w_{k-1}],$$

$$J(z_0) \leq \varepsilon + \sum_{k=1}^{\infty} [z_k^* (M+N) z_k + 2\operatorname{Re} z_k^* P z_{k-1}];$$

consequently,

$$\begin{aligned} J(w_0) + J(z_0) &\leq 2\varepsilon + \frac{1}{2} \sum_{k=1}^{\infty} [(w_k + z_k)^* (M+N) (w_k + z_k) + \\ &\quad + 2\operatorname{Re} (w_k + z_k)^* P (w_{k-1} + z_{k-1})] \\ &\quad + \frac{1}{2} \sum_{k=1}^{\infty} [(w_k - z_k)^* (M+N) (w_k - z_k) + \\ &\quad + 2\operatorname{Re} (w_k - z_k)^* P (w_{k-1} - z_{k-1})] \\ &\leq 2\varepsilon + \frac{1}{2} J(w_0 + z_0) + \frac{1}{2} J(w_0 - z_0). \end{aligned}$$

As  $\varepsilon > 0$  is arbitrary,  $2J(w_0) + 2J(z_0) \leq J(w_0 + z_0) + J(w_0 - z_0)$ ; by an obvious change of variables and quadratic homogeneity, the reverse inequality holds as well. Hence  $J^{1/2}(\cdot)$  satisfies the parallelogram law.

Define  $q: \mathbb{C}^n \times \mathbb{C}^n \rightarrow \mathbb{C}$  by

$$q(w, z) \equiv \frac{1}{4} [J(w+z) - J(w-z)] + \frac{i}{4} [J(w+iz) - J(w-iz)], \quad w, z \in \mathbb{C}^n.$$

We note that  $q$  is continuous with  $q(w, w) = J(w) \geq 0$ ,  $q(w, z) = \overline{q(z, w)}$ , and a standard argument [10] shows that

$$q(\alpha w + \hat{w}, z) = \alpha q(w, z) + q(\hat{w}, z) \quad \text{for all } w, \hat{w}, z \in \mathbb{C}^n, \quad \alpha \in \mathbb{C}.$$

Consequently, Riesz' representation theorem implies the existence of a unique Hermitian matrix  $Q$  such that  $z^* Q w = q(w, z) + z^* N w$ ,  $z, w \in \mathbb{C}^n$ .

Finally, we note that, for every  $u_1 \in \mathbb{C}^n$ ,

$$q(u_1, u_1) = J(u_1) = \sup_{\{u_k\}_{k=2}^{\infty}} \sum_{k=2}^{\infty} [u_k^* (M+N) u_k + 2 \operatorname{Re} u_k^* P u_{k-1}]$$

and therefore, for every  $u_0, u_1 \in \mathbb{C}^n$ ,

$$q(u_1, u_1) + u_1^* (M+N) u_1 + 2 \operatorname{Re} u_1^* P u_0 \leq J(u_0) = q(u_0, u_0);$$

consequently,

$$u_1^* (Q-N) u_1 + u_1^* (M+N) u_1 + 2 \operatorname{Re} u_1^* P u_0 \leq u_0^* (Q-N) u_0,$$

and we conclude that (i) implies (ii). ■

Through the use of this lemma we can provide existence criteria for Liapunov functions of the form (1.3) or (1.4) when equation (1.1) involves only one delay; i.e.,

$$\dot{y}(t) = Ay(t) + By(t-\tau), \quad t \geq 0, \quad (2.1)$$

with  $m = 1$ ,  $\tau_1 = \tau$ ,  $B_1 \equiv B$ . In this case the function (1.3) has the form

$$V(x) = y^* R y + \int_{-\tau}^0 v^*(\theta) Q v(\theta) d\theta, \quad x = (y, v) \in \chi, \quad (2.2)$$

for some  $n \times n$  Hermitian matrices  $R$  and  $Q$ . The following theorem gives necessary and sufficient condition for (2.2) to be a Liapunov function on  $\chi$  for (2.1).

Theorem 2.1: Given arbitrary  $n \times n$  positive-semidefinite Hermitian matrices  $C, D$ , the following two statements or equivalent:

- (i) There exist  $R = R^*$ ,  $Q = Q^*$ , such that  $V$  given by (2.2) is a nontrivial Liapunov function for (2.1) on  $\chi$  and

$$\dot{V}(x) \leq -y^* C y - v^*(-\tau) D v(-\tau), \quad x = (y, v) \in \chi.$$

- (ii) There exists  $R = R^* \neq 0$  such that

$$R A + A^* R + C + D + e^{i\phi} R B + e^{-i\phi} B^* R \leq 0 \quad \text{for all } \phi \in \mathbb{R}.$$

Proof: By direct computation of  $\dot{V}$  (see 1.7), we find that  $\dot{V}(x)$  satisfies the estimate of (i) if and only if the  $2n \times 2n$  Hermitian matrix

$$\begin{bmatrix} RA + A^*R + Q + C & BR \\ R^*B & D - Q \end{bmatrix}$$

is negative semidefinite; hence, Lemma 2.1 shows that (i) is equivalent to (ii). ■

This theorem leads to the following criteria for stability, asymptotic stability and instability of the trivial solution (hence, by linearity, of all solutions) of equation (2.1).

Theorem 2.2: Let there exist  $n \times n$  Hermitian matrices  $C, R$ , such that

$$RA + A^*R + C + e^{i\phi}RB + e^{-i\phi}B^*R \leq 0 \quad \text{for all } \phi \in \mathbb{R}. \quad (2.3)$$

Then, for all  $\tau \geq 0$ ,

- (i) If  $C \geq 0$  and  $R > 0$ , the trivial solution of (2.1) is stable.
- (ii) If  $C > 0$  and  $R > 0$ , the trivial solution of (2.1) is asymptotically stable;

- (iii) If  $C > 0$  and  $y_0^* R y_0 < 0$  for some  $y_0 \in \mathbb{C}^n$ ,  
the trivial solution of (2.1) is unstable.

Proof: By Lemma 2.1, condition (2.3) implies the existence of  $Q = Q^*$  such that

$$\begin{bmatrix} RA + A^*R + Q + C & RB \\ B^*R & -Q \end{bmatrix} \leq 0;$$

hence,  $Q \geq 0$ ,  $V(x) \geq y^* R y$ , and  $\dot{V}(x) \leq -y^* C y$  for the function  $V(x) = y^* R y + \int_{-\tau}^0 v^*(\phi) Q v(\phi) d\phi$ ,  $x = (y, v) \in \chi$ .

Using the semigroup notation of Section 1, we recall that  $C \geq 0$  implies  $V(S(t)x) \leq V(x)$ ,  $t \geq 0$ ,  $x \in \chi$ ; hence,  $V(x_0) \geq V(S(t)x_0) \geq y^*(t) R y(t)$ ,  $t \geq 0$ . Consequently, if  $R > 0$  and  $\|x_0\|_\chi \leq \delta$ , there exists  $c_0 \geq 1$  such that  $\|y(t)\|_\mathbb{C}^2 \leq c_0 \delta^2$ ,  $t \geq 0$ ; this implies that  $\|S(t)x_0\|_\chi^2 \leq c_0 \delta^2$  for all  $t \geq 0$ . It follows that (i) has been proved; moreover, for  $R > 0$ , every motion  $S(\cdot)x_0: \mathbb{R}^+ \rightarrow \chi$  has bounded positive orbit  $\gamma(x_0) \equiv \bigcup_{t \geq 0} S(t)x_0$ . It is well known that, for (1.2), bounded positive orbits are precompact [7]; hence, making the stronger assumption that  $C > 0$  and noting that the largest positive invariant set [11] in  $\{x \in \chi | \dot{V}(x) = 0\}$  is  $\{0\}$ , the Invariance Principle [11] yields the conclusion that  $S(t)x_0 \rightarrow 0$  as  $t \rightarrow \infty$ , for every  $x_0 \in \chi$ , which proves (ii).

Now considering (iii), with  $C > 0$  and  $y_0^* R y_0 < 0$  for some  $y_0 \in \mathbb{C}^n$ , the Invariance Principle shows that  $\gamma(x_0)$  is not precompact (hence, not bounded) for some  $x_0$  in each neighborhood of  $0 \in \chi$ ; hence,  $x = 0$  is unstable and the proof is complete. ■

We remark that if only the hypotheses of (i) are satisfied, rather than the stronger hypotheses of (ii), it may still be possible to employ the Invariance Principle [11] to conclude asymptotic stability rather than mere stability; one need only be able to show that the largest invariant set in  $\{x \in \chi | \dot{V}(x) = 0\}$  is  $\{0\}$ .

Let us now consider the function  $W$  of (1.4),

$$W(x) = \sup_{-\tau \leq \theta \leq 0} v^*(\theta) R v(\theta), \quad x = (y, v) \in \chi. \quad (2.4)$$

Recall that, for this function to be a Liapunov function for equation (2.1), it is required that

$$\operatorname{Re} z_0^* R (A z_0 + B z_1) \leq 0 \quad (2.5)$$

for all  $z_0, z_1 \in \mathbb{C}^n$  such that  $z_0^* R z_0 \geq z_1^* R z_1$ . By making the particular choice  $z_1 = e^{i\phi} z_0$  we note that a consequence of



condition (2.5) is that condition (2.3) must hold for  $C = 0$ . Hence existence of a Liapunov function of the form (2.4) implies the existence of a Liapunov function of the form (2.2), but not conversely unless, in (2.2), we have that  $Q = \alpha R$  for some  $\alpha > 0$ . It follows that the class of equations of the form (2.1) which admit "Razumikhin-type" Liapunov functions is a subset of the class that admits Liapunov functions of the form (2.2).

### III. The Multi-delay Case.

In this section we generalize the results presented in the previous section so as to encompass the case in which several delays occur, as in equation (1.1).

For this purpose, we first present a generalization of Lemma 2.1.

Lemma 3.1: Given  $n \times n$  complex matrices  $M, N_k, P_k$ , with  $M$  and  $N_k$  Hermitian,  $k = 1, 2, \dots, m$ , the following two statements are equivalent:

- (i) There exist Hermitian matrices  $M_k$ ,  $k = 1, 2, \dots, m$ , such that  $M \leq \sum_{k=1}^m M_k$  and, for each  $k = 1, 2, \dots, m$ ,  $M_k + N_k + e^{i\phi} P_k + e^{-i\phi} P_k^* \leq 0$  for all  $\phi \in \mathbb{R}$ .
- (ii) There exist Hermitian matrices  $Q_k$ ,  $k = 1, 2, \dots, m$ , such that

$$\begin{bmatrix} M + \sum_{k=1}^m Q_k & P_1 & P_2 & \dots & P_m \\ P_1^* & N_1 - Q_1 & 0 & \dots & 0 \\ P_2^* & 0 & N_2 - Q_2 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ P_m^* & 0 & 0 & \dots & N_m - Q_m \end{bmatrix} \leq 0.$$

Proof: It is clear that (ii) holds if and only if there exist Hermitian matrices  $H_k$ ,  $k = 1, 2, \dots, m$ , such that

$$\sum_{k=1}^m H_k \geq M + \sum_{k=1}^m Q_k \quad \text{and}$$

$$\begin{bmatrix} H_k & P_k \\ P_k^* & N_k - Q_k \end{bmatrix} \leq 0, \quad k = 1, 2, \dots, m.$$

Defining  $M_k \equiv H_k - Q_k$  and applying Lemma 1 for each  $k = 1, 2, \dots, m$ , it follows that (i) is equivalent to (ii). ■

As in the previous section, we will use this Lemma to provide existence criteria for Liapunov functions of the form (1.3) or (1.4) for equation (1.1),

$$\dot{y}(t) = Ay(t) + \sum_{k=1}^m B_k y(t - \tau_k), \quad t \geq 0. \quad (3.1)$$

We recall that we are interested in a function  $V$  having the form (1.3),

$$V(x) = y^* R y + \sum_{k=1}^m \int_{-\tau_k}^0 v^*(\theta) Q_k v(\theta) d\theta, \quad x = (y, v) \in X, \quad (3.2)$$

for some  $n \times n$  Hermitian matrices  $R$  and  $Q_k$ ,  $k = 1, \dots, m$ .

Theorem 3.1: Given arbitrary  $n \times n$  positive-semidefinite Hermitian matrices  $C, D_k$ ,  $k = 1, 2, \dots, m$ , the following two statements are equivalent:

- (i) There exist  $R = R^*$ ,  $Q_k = Q_k^*$ , such that  $V$  given by (3.2) is a nontrivial Liapunov function for (3.1) on  $X$  and

$$\dot{V}(x) \leq -y^*Cy - \sum_{k=1}^m v^*(-\tau_k)D_k v(-\tau_k), \quad x = (y, v) \in X.$$

- (ii) There exist  $R = R^* \neq 0$ ,  $M_k = M_k^*$ , such that  
 $RA + A^*R + C \leq \sum_{k=1}^m M_k$  and, for each  $k = 1, 2, \dots, m$ ,

$$M_k + D_k + e^{i\phi}RB_k + e^{-i\phi}B_k^*R \leq 0 \quad \text{for all } \phi \in \mathbb{R}.$$

The proof of this theorem follows immediately from application of Lemma 3.1, using the same arguments as in the proof of Theorem 2.1.

This result immediately leads to

Theorem 3.2: Let there exist  $n \times n$  Hermitian matrices  $C, M_k, R$  such that  $RA + A^*R + C \leq \sum_{k=1}^m M_k$  and, for each  $k = 1, 2, \dots, m$ ,

$$M_k + e^{i\phi}RB_k + e^{-i\phi}B_k^*R \leq 0 \quad \text{for all } \phi \in \mathbb{R}.$$

Then, for all  $\tau_k \geq 0$ ,  $k = 1, 2, \dots, m$ ,

- (i) If  $C \geq 0$  and  $R > 0$ , the trivial solution of (3.1) is stable.
- (ii) If  $C > 0$  and  $R > 0$ , the trivial solution of (3.1) is asymptotically stable.
- (iii) If  $C > 0$  and  $y_0^* R y_0 < 0$  for some  $y_0 \in \mathbb{C}^n$ , the trivial solution of (3.1) is unstable.

We omit the proof of this theorem since it parallels that of Theorem 2.2. As for Theorem 2.2, we remark that it may be possible to conclude asymptotic stability in certain applications satisfying the hypotheses of (i) but not those of (ii); this merely requires a more detailed application of the Invariance Principle [11].

The remarks made in Section II regarding the existence of a "Razumikhin-type" Liapunov function (i.e., of the form (1.4)) apply here as well; existence of this type of Liapunov function implies existence of a Liapunov function of the form (3.2), but the converse need not hold unless  $Q_k = \alpha_k R$ ,  $k = 1, 2, \dots, m$ , for some real  $\alpha_k > 0$ .



#### IV. A Converse Question on Asymptotic Stability.

If the hypotheses of conclusion (ii) of Theorem 3.2 are satisfied, then we are guaranteed asymptotic stability of all solutions of

$$\dot{y}(t) = Ay(t) + \sum_{k=1}^m B_k y(t-\tau_k), \quad t \geq 0, \quad (4.1)$$

irrespective of the values of the delays  $\tau_k$ ,  $k = 1, \dots, m$ . It follows that no function  $V$  of the form (1.3) will be a Liapunov function with  $R > 0$ ,  $\dot{V}(x) \leq -y^*Cy$ ,  $C > 0$ , unless equation (4.1) is asymptotically stable for all delays. Given this observation, and recalling the power of the Invariance Principle [11], it is natural to ask the following question: given that the trivial solution of (4.1) is asymptotically stable, irrespective of the delays  $\tau_k$ ,  $k = 1, \dots, m$ , does there necessarily exist a Liapunov function of the form (1.3) with  $R > 0$ ? This is essentially the question posed by Hale (see [7], p.108).

To investigate this question, let us recall a recent result [12] that characterizes those  $(m+1)$ -tuples of real matrices  $(A, B_1, \dots, B_m)$  such that the trivial solution of (4.1) will be asymptotically stable irrespective of the delays.

Theorem 4.1 [12]: The trivial solution of equation (4.1) is asymptotically stable for any set of  $\tau_k \geq 0$  if and only if

- (i) the real parts of the eigenvalues of  $A + \sum_{k=1}^m B_k$  are



negative, and

ii) for all  $\phi_k \in \mathbb{R}$ ,  $A + \sum_{k=1}^m e^{i\phi_k B_k}$  has no nonzero imaginary eigenvalue.

Thus, in the case of real matrices  $A, B_1, \dots, B_m$ , our question reduces to the following: do the hypotheses of Theorem 4.1 imply the existence of Hermitian matrices  $R > 0$ ,  $C \geq 0$ ,  $D_k \geq 0$  and  $M_k$ ,  $k = 1, \dots, m$ , such that statement (ii) in Theorem 3.1 is true?

This question has an affirmative answer in the scalar case (i.e.,  $A = a \in \mathbb{R}$ ,  $B_k = b_k \in \mathbb{R}$ ). Indeed the conditions of Theorem 4.1 become, in this case

$$a + \sum_{k=1}^m b_k < 0, \quad (4.2)$$

$$a + \sum_{k=1}^m |b_k| \leq 0. \quad (4.3)$$

Hence, letting  $R = 1$ ,  $C = 0$ ,  $D_k = 0$ , and  $M_k = -2|b_k|$ , it is immediately seen from (4.3) that all conditions of statement (ii) in Theorem 3.1 are satisfied, and, therefore, there exists a Liapunov function of the form

$$V(x) = |y|^2 + \sum_{k=1}^m \int_{-\tau_k}^0 Q_k |v(\theta)|^2 d\theta, \quad x = (y, v) \in X, \quad (4.4)$$

for some  $Q_1, Q_2, \dots, Q_m \in \mathbb{R}$ , such that  $\dot{V}(x) \leq 0$ . From the proof of Lemma 3.1, it is easily seen that the  $Q_k$  must satisfy

$$\begin{bmatrix} -2|b_k| + Q_k & b_k \\ b_k & -Q_k \end{bmatrix} \leq 0, \quad k = 1, 2, \dots, m;$$

hence,  $Q_k = |b_k|$  is the only possible choice, and we obtain

$$\dot{V}(x) = 2(a + \sum_{k=1}^m |b_k|) |y|^2 - \sum_{k=1}^m |b_k| [y - (\operatorname{sgn} b_k) v(-\tau_k)]^2$$

for all  $x = (y, v) \in X$ .

Although  $C \leq 0$  and result (ii) of Theorem 3.2 may not apply, the Invariance Principle [11] can be employed directly to show that, under condition (4.2), this Liapunov function does establish asymptotic stability.

The above question, however, has a negative answer at the vector level, as we can show through a counterexample. Indeed, in (4.1) let  $m = 1$  and

$$A = \begin{bmatrix} -1 & -1 \\ 1 & -1 \end{bmatrix}, \quad B_1 = \begin{bmatrix} \alpha & 0 \\ 0 & -\alpha \end{bmatrix},$$

for some  $\alpha \in \mathbb{R}$ . It is easily verified that the conditions of Theorem 4.1 are satisfied if and only if  $\alpha^2 < 2$ . On the other hand, it is not possible to satisfy condition (2.3) with any  $R = R^* > 0$ ,  $C = C^* \geq 0$ , when  $\alpha^2 > 1$ . Indeed, suppose

$$1 < \alpha^2 < 2 \quad \text{and}$$

$$R(A+e^{i\phi}B_1) + (A+e^{i\phi}B_1)^*R \leq -C \quad \text{for all } \phi \in \mathbb{R}, \quad (4.5)$$

where  $C = C^* \geq 0$  and

$$R = \begin{bmatrix} 1 & \bar{\beta} \\ \beta^* & \delta \end{bmatrix}, \quad \beta \in \mathbb{C}, \alpha \in \mathbb{R}, \delta > |\beta|^2.$$

Satisfaction of (4.5) implies that, for all  $\phi \in \mathbb{R}$ ,

$$\begin{bmatrix} -2+2\alpha\cos \phi + 2\operatorname{Re}\beta & -1-2\beta+\delta-2\alpha\beta\sin \phi \\ -1-2\bar{\beta}+\delta-2\alpha\bar{\beta}\sin \phi & -2\operatorname{Re}\beta-2\delta-2\delta\alpha\cos \phi \end{bmatrix} \leq 0.$$

For this to be so, it is necessary that

$$-2\delta(1+\alpha\cos \phi) \leq 2\operatorname{Re}\beta \leq 2(1-\alpha\cos \phi)$$

for all  $\phi \in \mathbb{R}$ , and this is impossible for  $1 < \alpha^2$ . Hence, for  $1 < \alpha^2 < 2$ , we have an example displaying asymptotic stability for all  $\tau \geq 0$ , but for which no nontrivial function of the form (1.3) is a Liapunov function.

# V. Additional Remarks.

In this section we present two additional results based on the use of a simple Liapunov function of the form (1.3). The first of these provides a sufficient condition for, and an estimate on, exponential decay of solutions. The second provides sufficient conditions for stability and asymptotic stability that are simpler than, but not as "necessary" as, those of Theorem 3.2.

In result (ii) of Theorem 3.2, and also (ii) of Theorem 2.2, we have concluded only asymptotic stability; without additional assumptions, this conclusion can be strengthened to exponential stability by employing yet another function, closely related to the function  $V$  of (1.3),

$$V_{\delta}(x) \equiv y^* R y + \sum_{k=1}^m \int_{-\tau_k}^0 e^{\delta(\theta+\tau_k)} v^*(\theta) Q_k v(\theta) d\theta, \quad x = (y, v) \in X, \quad (5.1)$$

where  $R = R^*$ ,  $Q_k = Q_k^*$ , and  $\delta$  is a positive real number. Computing  $\dot{V}_{\delta}: X \rightarrow \mathbb{R}$  according to (1.2), we obtain

$$\begin{aligned} \dot{V}_{\delta}(x) = & y^* (R A + A^* R + \sum_{k=1}^m e^{\delta \tau_k} Q_k) y + 2 \operatorname{Re} \sum_{k=1}^m y^* R B_k v(-\tau_k) \\ & - \sum_{k=1}^m v^*(-\tau_k) Q_k v(-\tau_k) \\ & - \delta \sum_{k=1}^m \int_{-\tau_k}^0 e^{\delta(\theta+\tau_k)} v^*(\theta) Q_k v(\theta) d\theta, \quad x = (y, v) \in X. \end{aligned} \quad (5.2)$$

Under the hypotheses of (ii) in Theorem 3.2 there exist  $C = C^* > 0$ ,  $R = R^* > 0$ ,  $Q_k = Q_k^*$  such that the Hermitian matrix

$$\begin{bmatrix} RA + A^*R + C + \sum_{k=1}^m Q_k & RB_1 & RB_2 & \dots & RB_m \\ B_1^*R & -Q_1 & 0 & \dots & 0 \\ B_2^*R & 0 & -Q_2 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ B_m^*R & 0 & 0 & \dots & -Q_m \end{bmatrix}$$

is negative-semidefinite. Consequently,  $Q_k \geq 0$  and the use of these matrices in  $V_\delta$  leads to

$$\begin{aligned} \dot{V}_\delta(x) &\leq -\delta V_\delta(x) + y^*(RA + A^*R + \delta R + \sum_{k=1}^m e^{+\delta \tau_k} Q_k)y + 2\operatorname{Re} \sum_{k=1}^m y^*RB_k v(-\tau_k) \\ &\quad - \sum_{k=1}^m v^*(-\tau_k) Q_k v(-\tau_k) \\ &\leq -\delta V_\delta(x) - y^*[C - \delta R - \sum_{k=1}^m (e^{\delta \tau_k} - 1)Q_k]y. \end{aligned}$$

Since  $C > 0$ , we may now choose a  $\delta > 0$  so small that

$\dot{V}_\delta(x) \leq -\delta V_\delta(x)$ , which leads to the estimate

$$V_\delta(S(t)x) \leq e^{-\delta t} V_\delta(x) \quad \text{for all } t \geq 0, x \in X. \quad (5.3)$$



Consequently,

$$y^*(t)Ry(t) \leq e^{-\delta t} \{v_0^*(0)Rv_0(0) + \sum_{k=1}^m e^{\delta \tau_k} \int_{-\tau_k}^0 v_0^*(\theta)Q_k v(\theta) d\theta\} \quad (5.4)$$

for all  $t \geq 0$ , implying the existence of  $c_0 \geq 1$  such that  $\|S(t)x_0\|_{\chi}^2 \leq c_0 e^{-\delta t} \|x_0\|_{\chi}^2$  for all  $t \geq 0$ ,  $x_0 \in \chi$ . It follows that the hypotheses of (ii) in Theorem 3.2 (hence, of (ii) in Theorem 2.2) actually imply exponential stability of the trivial solution, with exponent  $-\delta t/2$  such that  $\delta > 0$  and  $C \geq \delta R + \sum_{k=1}^m (e^{\delta \tau_k} - 1)Q_k$ ; hence,  $\delta$  depends on the delays  $\tau_k$ ,  $k = 1, 2, \dots, m$ .

Turning to the second remark we wish to make, we refer again to the function  $V$  of (1.3), recalling that  $V$  is a Liapunov function for (1.2) if and only if (1.7) holds; i.e.,

$$\begin{bmatrix} RA + A^*R + \sum_{k=1}^m Q_k & RB_1 & RB_2 & \dots & RB_m \\ B_1^*R & -Q_1 & 0 & \dots & 0 \\ B_2^*R & 0 & -Q_2 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ B_m^*R & 0 & 0 & \dots & -Q_m \end{bmatrix} \leq 0 \quad (5.5)$$



for some  $R = R^*$ ,  $Q_k = Q_k^*$ ,  $k = 1, 2, \dots, m$ . Theorem 3.1 shows this condition to be satisfied if and only if statement (ii) of that theorem is true for  $C \equiv 0$ ,  $D_k \equiv 0$ ,  $k = 1, 2, \dots, m$ . Unfortunately, this statement may be difficult to verify and does not suggest an explicit construction for the matrices  $Q_k$ . In certain applications it may be desirable instead to simply "guess" a plausible form for  $Q_k = Q_k^*$ , and then search for some  $R = R^*$  such that (5.5) holds. We will now describe a simple technique of this type.

Condition (5.5) implies that  $Q_k \geq 0$  with null space no larger than that of  $RB_k$ . Hence, let us suppose  $Q_k$  to be of the form  $Q_k = B_k^* R Z_k R B_k$  for some  $Z_k = Z_k^* > 0$ ,  $k = 1, 2, \dots, m$ . Then (5.5) will be satisfied if there exists  $R = R^*$ ,  $Z_k = Z_k^* > 0$ , such that

$$RA + A^*R \leq - \sum_{k=1}^m (B_k^* R Z_k R B_k + Y_k) \quad (5.6)$$

where  $Y_k$  is a Hermitian matrix that inverts  $Z_k$  on the range of  $RB_k$  ( $Y_k = Z_k^{-1}$  if  $\det |RB_k| \neq 0$ ). In fact, a careful review of the proofs of Lemma 2.1 and Lemma 3.1 reveals that if (5.5) can be satisfied by any  $R = R^* \neq 0$ ,  $Q_k = Q_k^*$ , then it can be satisfied by some  $R = R^* \neq 0$ ,  $Q_k = Q_k^*$ , with  $Q_k$  of the form we have suggested. Hence, satisfaction of (5.6) by some  $R = R^* \neq 0$ ,  $Z_k = Z_k^* > 0$ , is necessary and sufficient for the existence of any non-trivial Liapunov function of the form (1.3) for equation (1.2).

Condition (5.6) certainly will be satisfied if there exists  $R = R^* \geq 0$  and real numbers  $\beta_k > 0$  such that

$$RA + A^*R + \sum_{k=1}^m (\beta_k B_k^* R B_k + \beta_k^{-1} R) \leq 0. \quad (5.7)$$

Although much more restrictive than our necessary and sufficient conditions, the comparatively simple condition (5.7) is certainly sufficient to ensure that  $V$  is a Liapunov function on  $\chi$  for (1.3) with  $Q_k \equiv \beta_k B_k^* R B_k$ , and that the trivial solution is stable if  $R > 0$ ; moreover,  $\dot{V}$  is given by

$$\begin{aligned} \dot{V}(x) = & y^* [RA + A^*R + \sum_{k=1}^m (\beta_k B_k^* R B_k + \beta_k^{-1} R)] y \\ & - \beta_k^{-1} [y - \beta_k B_k v(-\tau_k)]^* R [y - \beta_k B_k v(-\tau_k)], \end{aligned}$$

$$x = (y, v) \in \chi.$$

Hence, under (5.7), if  $R > 0$ , it is possible to use the Invariance Principle [11] to provide conditions sufficient for asymptotic stability; for example, it is easily found that the trivial solution is asymptotically stable if  $R = R^* > 0$  and  $\beta_k > 0$ ,  $k = 1, 2, \dots, m$ , are such that inequality (5.7) holds strictly.

## VI.

REFERENCES

- [1] Yoshizawa, T., Stability Theory by Liapunov's Second Method, Math. Soc. Japan, 1966.
- [2] Yoshizawa, T., Stability Theory and the Existence of Periodic Solutions and Almost Periodic Solutions, Appl. Math. Sciences, Vol. 14, Springer Verlag, 1975.
- [3] Razumikhin, B.S., Application of Liapunov's Method to problems in the stability of systems with delays, Autom. i Telemekh, 21, (1960), 740-749.
- [4] Krasovskii, M., Stability of Motion, Stanford Univ. Press, 1963.
- [5] Driver, R.D., Linear differential systems with small delays, J. Diff. Equations, 21, (1976) 148-166.
- [6] Hale, J.K., Functional Differential Equations, Appl. Math. Sciences, Vol. 3, Springer-Verlag, 1971.
- [7] Hale, J.K., Theory of Functional Differential Equations, Applied Math. Sciences, Vol. 3, Springer-Verlag, 1977.
- [8] Infante, E.F. and J.A. Walker, A Liapunov functional for a scalar differential difference equation, Proc. Royal Soc. of Edinburgh, 79A, 1977, 307-316.
- [9] DeBona Castelan, W. and E.F. Infante, A Liapunov functional for a matrix difference differential equation, J. Diff. Equations, 29, 1978, 439-451.
- [10] Yoshida, K., Functional Analysis, 4th Edition, Springer-Verlag, 1974.
- [11] Hale, J.K., Dynamical Systems and Stability, J. Math. Anal. and Appl., 26, 1969, 39-59.
- [12] Hale, J.K., E. F. Infante and F.P. Tsen, Stability in delay equations for all delays, Center for Dynamical Systems Report, Brown University, 1979.